

Problem III.3 . . . randomly you get further

5 points; průměr 4,23;

řešilo 86 studentů

In the microworld of cells, there are two types of transport: transport by free diffusion, also known as Brownian motion where the motion uses the energy of the environment directly. The second type, so-called active transport, requires, among other things, a motor protein moving at a constant speed along the cytoskeletal filament. Consider the typical value of the diffusion constant $D \approx 10^{-9} \text{ cm}^2 \cdot \text{s}^{-1}$ and the rate of active transport speed $u \approx 10^{-6} \text{ m} \cdot \text{s}^{-1}$. For which distances is the Brownian motion more time efficient than the active transport? Assume that the transport is happening in only one direction.

Marek J. read Sekimoto.

The mean distance travelled for a free Brownian particle in 1D is given by

$$\langle x \rangle = \sqrt{2Dt},$$

where t is the time and $\langle \rangle$ denotes the average over different trajectories. A diffusive motion mode is preferred for short times and/or distances since its cause is the surrounding environment. Specifically, the motion is generated by the collisions of the molecules of the environment with the Brownian particle. For longer distances, therefore longer times, randomness can no longer be relied on. The mean velocity $\langle x \rangle / t = \sqrt{2D/t}$ decreases with time. The distance l for which both transport methods are on average equally fast (it takes the same time τ to travel l), is determined as follows

$$\tau = \frac{l^2}{2D} = \frac{l}{u},$$

using the values from the problem statement, we have $l = 2D/u \approx 2 \cdot 10^{-7} \text{ m}$. We see that diffusive motion can be preferable to active transport for distances smaller than the order of hundreds of nanometers. Nature indeed takes advantage of this. For example, the distance of roughly 50 nm between synaptic membranes is overcome by neurotransmitters via diffusion. On the other hand, distances in cells, of the order of micrometres, are typically overcome by molecular motors.

Marek Jankola
marekj@fykos.org

FYKOS is organized by students of Faculty of Mathematics and Physics of Charles University. It's part of Media Communications and PR Office and is supported by Institute of Theoretical Physics of MFF UK, his employees and The Union of Czech Mathematicians and Physicists. The realization of this project was supported by Ministry of Education, Youth and Sports of the Czech Republic.

This work is licensed under Creative Commons Attribution-Share Alike 3.0 Unported. To view a copy of the license, visit <https://creativecommons.org/licenses/by-sa/3.0/>.